

AD-A049 443

NAVAL RESEARCH LAB WASHINGTON D C  
TRANSIENT RESPONSE OF TWO FLUID-COUPLED SPHERICAL ELASTIC SHELL--ETC(U)  
DEC 77 H HUANG

F/G 20/11

JNCLASSIFIED

NRL-8185

SBIE-AD-E000 099

NL

1 OF 1  
AD  
A049443



END  
DATE  
FILMED  
3 - 78  
DOC

AD A049443

AD No. \_\_\_\_\_  
DDC FILE COPY

ade 000099  
NRL Report 8185

12  
£

# Transient Response of Two Fluid-Coupled Spherical Elastic Shells to an Incident Pressure Pulse

H. HUANG

*Ocean Technology Division*

December 23, 1977

This research was sponsored by the Defense Nuclear Agency under  
Subtask 99QAXSF502, work unit 10, work unit title Double Hull Response Evaluation



DDC  
RECEIVED  
FEB 6 1978  
B

NAVAL RESEARCH LABORATORY  
Washington, D.C.

Approved for public release: distribution unlimited.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER NRL 8185 ✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER	
4. TITLE (and Subtitle) TRANSIENT RESPONSE OF TWO FLUID-COUPLED SPHERICAL ELASTIC SHELLS TO AN INCIDENT PRESSURE PULSE.		5. TYPE OF REPORT & PERIOD COVERED Interim report on a continuing NRL Problem	
7. AUTHOR(s) H. Huang		6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Research Laboratory Washington, D. C. 20375 ✓		8. CONTRACT OR GRANT NUMBER(s)	
11. CONTROLLING OFFICE NAME AND ADDRESS Defense Nuclear Agency Washington, D.C. 20305		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NRL Problem MF02-12B DNA Subtask 99QAXSF502, Work Unit 10	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (12) 15p.		12. REPORT DATE 23 December 1977	
		13. NUMBER OF PAGES 28	
		15. SECURITY CLASS. (of this report) UNCLASSIFIED	
16. DISTRIBUTION STATEMENT (of this Report) (18) SBIE (19) AD-E000 099 Approved for public release; distribution unlimited.		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		DDC RECEIVED FEB 6 1978 RECEIVED B	
18. SUPPLEMENTARY NOTES This research was sponsored by the Defense Nuclear Agency under Subtask 99QAXSF502, work unit 10, work unit title Double Hull Response Evaluation			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Transient analysis Shock wave loading Fluid-structure interaction Double hull			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The transient response of a system of two concentric spherical elastic shells coupled by an ideal fluid and impinged by an incident plan pressure pulse is analyzed. The classical techniques of separation of variables and Laplace transforms are employed for solving the wave equations governing the fluid motions and the shell equations of motion. A scheme of iterative convolution was devised for the inversion of the Laplace transforms that facilitates the calculation of accurate transient solutions of the response of the shells. A sample calculation of shell responses was performed and results are compared to the case in which the outer shell is absent. This set of results demonstrates that a thin outer shell tends to be transparent to the incident pulse.			

DD FORM 1473  
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE  
5/N 0102-LF-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

251 950

LB

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

[Large empty rectangular box for content]



## CONTENTS

INTRODUCTION .....	1
Description of the Problem .....	2
Solutions in the Laplace Transform Domain .....	4
The Inversion of the Transformed Solutions .....	6
Results and Discussions .....	8
ACKNOWLEDGMENT .....	11
REFERENCES .....	12

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION _____	
BY _____	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	

## **TRANSIENT RESPONSE OF TWO FLUID-COUPLED SPHERICAL ELASTIC SHELLS TO AN INCIDENT PRESSURE PULSE**

### **INTRODUCTION**

An underwater weak shock wave sufficiently far away from its generating explosion source is often treated as an acoustic pulse. There is prolific literature on the studies of the transient interaction among such pressure waves and single elastic shells of simple shapes [1,2]. The results not only reveal many of the essential physical phenomena involved in the interaction problem but also are quite useful for the verification of approximation methods for predicting the underwater explosion response of submerged structures surrounded by an exterior fluid medium of infinite extent [2-5]. In the present endeavor, the transient response of a system of two fluid coupled concentric spherical elastic shells impinged by an external incident plane shock wave is analyzed. This purports to gain physical insight in the response as well as to provide a data base for the development of general numerical methods for predicting the underwater explosion response of fluid coupled shell systems such as the double hull section of a submarine. The problem with the spherical geometry permits the separation of variables in the wave equation governing the fluid motion and the shell equations of motion. The use of Laplace transforms then facilitates the calculation of satisfactory transient solutions of the response of the shells.

A paper appears in the Russian literature [6] independently dealing with exactly the same problem. The results obtained, however, only pertain to the point-symmetric and the translational motions of the shells and these two terms of the series solution are insufficient for the description of the complete shell response. It would also seem that the scheme of dual Volterra integral equations for the calculation of the inverse Laplace transform used in Ref. 6 is unnecessarily complex and numerically inefficient. It can be readily shown that only a single and simpler integral equation is needed if this scheme is used.

In this report, a simple, straightforward method is devised for the inverse Laplace transform, and the asymptotic behaviors of the shells are analytically discussed. An example calculation is also carried out using eight terms of the series solution for adequate convergence of shell displacements, velocities, and stresses. Time histories of the transient response of the interior shell are presented. This example demonstrates that a thin exterior shell tends to be transparent to the incident pulse.

## Description of the Problem

Figure 1 sketches the fluid-coupled spherical shell system and the incident plane pressure wave. The fluid surrounding the outer shell and that between the two shells are considered to be ideal compressible fluids in linear wave motions and can be characterized by their unperturbed mass densities and sound speeds, i.e., by  $(\rho^e, c^e)$  and  $(\rho, c)$ , respectively. The shells are initially concentric. In this study, the strength of the incident wave is sufficiently weak such that the shell deflections are elastic and small and the deviation from the concentricity remains negligible for the time duration of interest. The mass densities, Young's moduli and Poisson's ratios of the outer and inner shells are  $(\rho_s^e, E^e, \nu^e)$  and  $(\rho_s, E, \nu)$ , respectively. The middle surface radii and thicknesses of the outer and inner shells are  $(a^e, h^e)$  and  $(a, h)$ , respectively. The  $\Phi$ -coordinate of the spherical system  $(r, \theta, \Phi)$  is not shown in Fig. 1 since it is not needed due to the symmetry of the problem. The origin  $O$  coincides with the unperturbed center of the shells.




Fig. 1 - Geometry of the problem

The deflections of the inner shell in the  $r$ - and  $\theta$ -direction, normalized with respect to the outer shell radius  $a^e$ , are denoted by  $w$  and  $u$  respectively and those of the outer shell by  $w^e$  and  $u^e$  respectively. The total pressure field exterior to the outer shell is denoted by  $p^e(r, \theta, t)$  and that between the shells by  $p(r, \theta, t)$  where  $t$  designates time. The following dimensionless parameters will be used in the mathematical formulation:

$$\begin{aligned}
 R &= r/a^e, \quad T = c^e t/a^e, \quad \zeta = a/a^e, \quad \eta = c_r(1 - \zeta), \\
 c_r &= c^e/c, \quad \rho_r = \rho/\rho^e, \quad M = \rho^e a/(\rho_s h), \quad M^e = \rho^e a^e/(\rho_s^e h^e), \\
 \Pi &= p/[\rho^e (c^e)^2], \quad \Pi^e = p^e/[\rho^e (c^e)^2], \quad I = \frac{1}{12} (h/a)^2, \quad I^e = \frac{1}{12} (h^e/a^e)^2, \\
 C^2 &= E/[\rho_s (1 - \nu) (c^e)^2], \quad C_e^2 = E^e/[\rho_s^e (1 - \nu^e) (c^e)^2], \\
 \alpha_0 &= \alpha_0^e = \mu_0 = \mu_0^e = 0, \quad \lambda_0 = 2C^2, \quad \lambda_0^e = 2C_e^2, \\
 \alpha_m &= (1 + I)C^2 [m(m+1) - (1 - \nu)]/(1 + \nu), \\
 \alpha_m^e &= (1 + I^e)C_e^2 [m(m+1) - (1 - \nu^e)]/(1 + \nu^e), \\
 \lambda_m &= C^2 [2 + [1 + (m^2 + m + 1)I]/[m(m+1) - (1 - \nu)]/(1 + \nu)], \\
 \lambda_m^e &= C_e^2 [2 + [1 + (m^2 + m + 1)I^e]/[m(m+1) - (1 - \nu^e)]/(1 + \nu^e)], \\
 \mu_m &= C^4 (m^2 + m - 2) [1 - \nu + [m(m+1) - (1 - \nu)] \times \\
 &\quad [m(m+1) - (1 + \nu)]/[1 - \nu)]/(1 - \nu) \\
 \mu_m^e &= C_e^4 (m^2 + m - 2) [1 - \nu^e + [m(m+1) - (1 - \nu^e)] \times \\
 &\quad [m(m+1) - (1 + \nu^e)]/[1 - \nu^e)]/(1 - \nu^e) \\
 m &= 1, 2, 3, \dots
 \end{aligned} \tag{1}$$



$\Pi^e$  and  $\Pi$  satisfy the wave equations

$$\nabla^2 \Pi^e = \frac{\partial^2 \Pi^e}{\partial T^2} \quad (2)$$

and

$$\nabla^2 \Pi = c_r^2 \frac{\partial^2 \Pi}{\partial T^2} \quad (3)$$

respectively, where  $\nabla^2$  is the Laplacian operator. The boundary conditions of the problem are that  $\Pi^e$  satisfies the radiation condition at far field and that

$$\frac{\partial \Pi}{\partial R} = \frac{\partial \Pi^e}{\partial R} = -\frac{\partial^2 w^e}{\partial T^2} \text{ at } R = 1 \quad (4)$$

and

$$\frac{\partial \Pi}{\partial R} = -\rho_r \frac{\partial^2 w}{\partial T^2} \text{ at } R = \zeta. \quad (5)$$

All quantities except the incident pressure field have quiescent initial conditions.

A Laplace transform pair is defined as

$$\begin{aligned} \bar{w}(\theta, s) &= \int_0^\infty w(\theta, T) e^{-sT} dT \\ w(\theta, T) &= \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} \bar{w}(\theta, s) e^{sT} ds \end{aligned} \quad (6)$$

where  $\gamma$  lies to the right of all singularities of  $\bar{w}$  in the complex  $s$ -plane and  $i = (-1)^{1/2}$ .

Due to the spherical geometry of the problem, the solutions can be expanded in terms of series of Legendre polynomials as the following.

$$\begin{aligned} \Pi(R, \theta, T) &= \sum_{m=0}^{\infty} \Pi_m(R, T) P_m(\cos \theta) \\ \Pi^e(R, \theta, T) &= \sum_{m=0}^{\infty} \Pi_m^e(R, T) P_m(\cos \theta) \\ w(\theta, T) &= \sum_{m=0}^{\infty} w_m(T) P_m(\cos \theta) \\ w^e(\theta, T) &= \sum_{m=0}^{\infty} w_m^e(T) P_m(\cos \theta) \\ u(\theta, T) &= \sum_{m=1}^{\infty} u_m(T) \frac{dP_m(\cos \theta)}{d\theta} \\ u^e(\theta, T) &= \sum_{m=1}^{\infty} u_m^e(T) \frac{dP_m(\cos \theta)}{d\theta} \end{aligned} \quad (7)$$

where  $P_m$  is the Legendre polynomial of the first kind and  $m$ th degree.



In the Laplace transform domain, the equations of motion of the elastic shells are [7]

$$\begin{aligned}\bar{w}_m &= \frac{-M\zeta\Pi_m(\zeta, s)(\zeta^2s^2 + \alpha_m)}{\zeta^4s^4 + \lambda_m\zeta^2s^2 + \mu_m} \\ \bar{w}_m^e &= \frac{-M^e[\Pi_m^e(1, s) - \Pi_m(1, s)](s^2 + \alpha_m^e)}{s^4 + \lambda_m^es^2 + \mu_m^e} \\ m &= 0, 1, 2, \dots\end{aligned}\quad (8)$$

$$\begin{aligned}(\zeta^2s^2 + \alpha_m)\bar{u}_m &= \left[C^2 + \frac{I}{1+I}\alpha_m\right]\bar{w}_m \\ (s^2 + \alpha_m^e)\bar{u}_m^e &= \left[C_e^2 + \frac{I^e}{1+I^e}\alpha_m^e\right]\bar{w}_m^e \\ m &= 1, 2, 3, \dots\end{aligned}\quad (9)$$

It should be noted that the solution method developed here is applicable for any linear elastic theory for the spherical shells. The choice of the version in Eqs. (8) and (9) is for the comparison of results previously obtained in Ref. 7.

The total pressure field exterior to the outer shell  $\Pi^e$  consists of the pressure due to the incident wave and those due to scattering and radiation by the outer shell. An arbitrary incident plane pressure wave impinging the vertex of the outer shell ( $R = 1, \theta = 0$ ) at  $T = 0$  can be expressed by the following series [7]

$$\Pi^i(R, \theta, s) = f(s)e^{-s} \sum_{m=0}^{\infty} (2m+1)i_m(Rs)P_m(\cos\theta), \quad (10)$$

where  $f(s)$  is the Laplace transform of the time characteristics of the incident wave and  $i_m(Rs)$  is the abbreviated notation for the modified spherical Bessel function of the first kind  $[\pi/(2Rs)]^{1/2}I_{m+1/2}(Rs)$  [8].

### Solutions in the Laplace Transform Domain

It can be shown that the solutions to the system of Eqs. (2) through (10) are:

$$\begin{aligned}\bar{\Pi}_m^e(R, s) &= (2m+1) \frac{f(s)e^{-s}}{k_m'(s)} [i_m(Rs)k_m'(s) - k_m(Rs)i_m'(s)] \\ &\quad - \frac{s\bar{w}_m^ek_m'(Rs)}{k_m'(s)}\end{aligned}\quad (11)$$

$$\begin{aligned}\bar{\Pi}_m(R, s) &= \frac{\rho_r s}{c_r [i_m'(c_r\zeta s)k_m'(c_r s) - i_m'(c_r s)k_m'(c_r\zeta s)]} \\ &\quad \times \{ [\bar{w}_m^ek_m'(c_r\zeta s) - \bar{w}_mk_m'(c_r s)]i_m(c_r Rs) \\ &\quad - [\bar{w}_m^ei_m'(c_r\zeta s) - \bar{w}_mi_m'(c_r s)]k_m(c_r Rs) \}\end{aligned}\quad (12)$$

$$\begin{aligned} \bar{w}_m^e = & -\frac{(2m+1)}{\Delta_m(s)} \frac{\pi e^{-s}}{2s^2} f(s) M^e (s^2 + \alpha_m^e) \{ (\zeta^4 s^4 + \lambda_m \zeta^2 s^2 + \mu_m) \\ & \times [k_m'(c_r s) i_m'(c_r \zeta s) - i_m'(c_r \zeta s) k_m'(c_r s)] + \rho_r M \frac{\zeta s}{c_r} (\zeta^2 s^2 + \alpha_m) \\ & \times [i_m'(c_r s) k_m(c_r \zeta s) - i_m(c_r \zeta s) k_m'(c_r s)] \} \end{aligned} \quad (13)$$

$$\bar{w}_m = -\frac{(2m+1)}{\Delta_m(s)} \frac{\pi e^{-s}}{2s^2} f(s) \frac{\pi}{2c_r^3 \zeta s} \rho_r M M^e (\zeta^2 s^2 + \alpha_m) (s^2 + \alpha_m^e). \quad (14)$$

In the above equations  $k_m(s)$  is the abbreviated notation for the modified spherical Bessel function of the third kind,  $[\pi/2s]^{1/2} K_{m+1/2}(s)$ . The prime denotes differentiation of the Bessel functions with respect to their arguments, and

$$\begin{aligned} \Delta_m(s) = & \left\{ (\zeta^4 s^4 + \lambda_m \zeta^2 s^2 + \mu_m) [k_m'(c_r s) i_m'(c_r \zeta s) - k_m'(c_r \zeta s) i_m'(c_r s)] \right. \\ & \left. + \rho_r M \frac{\zeta s}{c_r} (\zeta^2 s^2 + \alpha_m) [i_m'(c_r s) k_m(c_r \zeta s) - k_m'(c_r s) i_m(c_r \zeta s)] \right\} \\ & \times [k_m'(s) (s^4 + \lambda_m^e s^2 + \mu_m^e) - M^e s k_m(s) (s^2 + \alpha_m^e)] \\ & - \frac{\rho_r}{c_r} M^e s (s^2 + \alpha_m^e) (\zeta^4 s^4 + \lambda_m \zeta^2 s^2 + \mu_m) k_m'(s) \\ & \times [k_m'(c_r \zeta s) i_m'(c_r s) - i_m'(c_r \zeta s) k_m(c_r s)] \\ & + \left( \frac{\rho_r}{c_r} \right)^2 M M^e \zeta s^2 (s^2 + \alpha_m^e) (\zeta^2 s^2 + \alpha_m) k_m'(s) \\ & \times [i_m'(c_r s) k_m(c_r \zeta s) - i_m(c_r \zeta s) k_m'(c_r s)] \end{aligned} \quad (15)$$

With use of the Tauber's theorem of Laplace transforms [9], some of the asymptotic behaviors of the shell responses at late time can readily be revealed from Eqs. (13) and (14). Specifically, for the case where the incident wave  $\Pi'$  is a unit step wave, i.e.,  $f(s) = 1/s$ ,

$$w_0^e(T) = \frac{-2C^2 c_r^2 (1 - \zeta^3) M^e + 3\zeta^3 \rho_r M M^e}{4C^2 C_e^2 c_r^2 (1 - \zeta^3) + 6\rho_r M \zeta^3 C_e^2 + 6\rho_r M^e C^2} \quad (16)$$

and

$$w_0(T) = \frac{-3\zeta \rho_r M M^3}{4C^2 C_e^2 c_r^2 (1 - \zeta^3) + 6\rho_r M \zeta^3 C_e^2 + 6\rho_r M^e C^2}. \quad (17)$$

These shell deflections occur long after the incident wave has engulfed the outer shell and can also be found by static analysis.

Again, for the unit step incidence case,

$$\begin{aligned} \dot{w}_1^e(T) = & \frac{-3M^e [(1+2\zeta^2)\rho_r M + 6(1-\zeta^3)]}{6\rho_r M (1+2\zeta^3) + 6M^e [(1-\zeta^3) + (2+\zeta^3)\rho_r] + \rho_r M M^e [1+2\zeta^3+2\rho_r(1-\zeta^3)] + 36(1-\zeta^3)} \end{aligned} \quad (18)$$

and

$$\dot{w}_1(T) = \frac{-9\rho_r MM^e}{6\rho_r M(1+2\zeta^3) + 6M^e[(1-\zeta^3) + (2+\zeta^3)\rho_r] + \rho_r MM^e[1+2\zeta^3+2\rho_r(1-\zeta^3)] + 36(1-\zeta^3)} \quad (19)$$

where the dot denotes differentiation with respect to  $T$ . These are late time translational velocities of the shells in the direction of propagation of the incident wave. It can be seen that the only condition under which  $\dot{w}_1 = \dot{w}_1^e$  at late time is  $\rho_r M = 3$ , i.e., when the inner shell is neutrally buoyant in the interior fluid. Otherwise, they are not equal to each other. Therefore, when interpreting the results of the present problem for large values of  $T$ , the relative positions of the shell must also be examined. If, at some time, the deviation from concentricity becomes excessive, the results of analysis thereafter are no longer physically meaningful.

Formulae (16) through (19) are convenient for parametric studies of the effects of various shell properties and fluid properties on the symmetric and translational responses of the shells, and they are also quite useful for providing asymptotic checks for the numerical calculations.

#### The Inversion of the Transformed Solutions

The spherical Bessel functions can be expressed in finite series of elementary functions [8]. If this property is used, the transformed solutions, e.g., Eq. (14), can be rearranged in the following form:

$$\bar{w}_m(s) = \frac{\Gamma_m(s)}{A_m(s)e^{\eta s} - B_m(s)e^{-\eta s}}, \quad (20)$$

where

$$\begin{aligned} \Gamma_m(s) &= -2(-1)^{m+1}(2m+1)(\rho_r/c_r)f(s)MM^e(\zeta s)^{m+1}(c_r s)^{2m+2}(s^2 + \alpha_m^e)(\zeta^2 s^2 + \alpha_m) \\ A_m(s) &= \{X_m(s)X_m(-c_r s)(s^4 + \alpha_m^e s^2 + \mu_m^e) + M^e s^2(s^2 + \alpha_m^e)[Y_m(s)X_m(-c_r s) \\ &\quad - \rho_r Y_m(-c_r s)X_m(s)]\}[X_m(c_r \zeta s)(\zeta^4 s^4 + \lambda_m \zeta^2 s^2 + \mu_m) \\ &\quad + \rho_r M \zeta^2 s^2(\zeta^2 s^2 + \alpha_m)Y_m(c_r \zeta s)] \\ B_m(s) &= \{X_m(s)X_m(c_r s)(s^4 + \lambda_m^e s^2 + \mu_m^e) + M^e s^2(s^2 + \alpha_m^e)[Y_m(s)X_m(c_r s) \\ &\quad - \rho_r Y_m(c_r s)X_m(s)]\}[X_m(-c_r \zeta s)(\zeta^4 s^4 + \lambda_m \zeta^2 s^2 + \mu_m) \\ &\quad + \rho_r M \zeta^2 s^2(\zeta^2 s^2 + \alpha_m)Y_m(-c_r \zeta s)] \\ X_m(s) &= \sum_{k=0}^{m+1} (m+3/2, k)2^{-k} s^{m+1-k} - m \sum_{k=0}^m (m+1/2, k)2^{-k} s^{m-k} \\ Y_m(s) &= \sum_{k=0}^m (m+1/2, k)2^{-k} s^{m-k} \\ (m+1/2, k) &= \frac{(m+k)!}{k!(m-k)!}. \end{aligned} \quad (21)$$



Both  $A_m$  and  $B_m$  are  $(3m + 11)$ th order real polynomials of  $s$  and, for the plane incidence case,  $\Gamma_m$  is a  $(3m + 6)$ th order real polynomial.

Equations of the form of Eq. (20) appear in many applied problems such as wave propagation in layered media, transmission lines, etc. The exponential factors in the denominator signify the reflections of waves, in the present case, to and from  $R = \zeta$  and  $R = 1$ . The dimensionless time for a disturbance to traverse the distance  $(1 - \zeta)$  is  $\eta$ . There is a variety of schemes for calculating the inverse Laplace transform of Eq. (20). One of the standardized techniques is the D'Alembert expansion [10] by which Eq. (20) can be rewritten as

$$\bar{w}_m(s) = \frac{\Gamma_m(s)e^{-\eta s}}{A_m(s)} \left\{ 1 + \frac{B_m}{A_m} e^{-2\eta s} + \left(\frac{B_m}{A_m}\right)^2 e^{-4\eta s} + \left(\frac{B_m}{A_m}\right)^3 e^{-6\eta s} + \dots \right. \\ \left. + \left(\frac{B_m}{A_m}\right)^n e^{-2n\eta s} + \dots \right\} \\ n = 1, 2, 3, \dots \quad (22)$$

Since  $A_m$  is a real polynomial of  $s$ , its roots can be accurately computed by well-established numerical procedures and the inversion of every individual term on the right-hand side of Eq. (22) can then be obtained by the method of residue. The calculation of residues, however, would be rather clumsy for terms with  $A_m$  of high power. The following scheme utilizing the convolution theorem is devised to circumvent this. From Eq. (22),

$$w_m(T) = w_m^0(T - \eta)H(T - \eta) + w_m^1(T - 3\eta) + w_m^2(T - 5\eta)H(T - 5\eta) \\ + \dots + w_m^n[T - (2n + 1)\eta]H[T - (2n + 1)\eta] + \dots \quad (23)$$

where  $H$  is the Heaviside step function and

$$w_m^1(T) = w_m^0(T) - \int_0^T G_m(T - \tau)w_m^0(\tau)d\tau, \\ w_m^2(T) = w_m^1(T) - \int_0^T G_m(T - \tau)w_m^1(\tau)d\tau, \\ \vdots \\ w_m^n(T) = w_m^{n-1}(T) - \int_0^T G_m(T - \tau)w_m^{n-1}(\tau)d\tau. \quad (24)$$

In Eq. (24),  $w_m^0(T)$  and  $G_m(T)$  are the inverse transforms of  $\Gamma_m(s)/A_m(s)$  and  $[A_m(s) - B_m(s)]/A_m(s)$  respectively. They are to be first accurately obtained by the method of residue. This is quite practical with the current computing technology for up to moderate values of  $m$ . For large  $m$  asymptotic expansions for the spherical Bessel functions can be used and  $A_m$  and  $B_m$  will assume simpler forms. For the calculations of shell responses in the present problem, large  $m$  terms are not needed. The successive convolutions required in Eq. (24) can be conveniently programmed and carried out in a modern digital computer. Since  $w_m^0(T)$  and  $G_m(T)$  are composed of terms formed by an exponential multiplied by a trigonometrical function, Trauboth's fast convolution integration algorithm [11], which only requires about the same number of computation steps as for ordinary integration, is used here.



It can be seen from Eqs. (23) and (24) that each successive integration advances the solution time by  $2\eta$  and the number of successive integrations required depends on  $T$  and  $\eta$ . The control of numerical accuracy lies in finding the zeros of  $A_m$  and the subsequent numerical integrations, and both are well established numerical techniques. Other quantities such as  $w_m^e(T)$  can be found by the same procedure or calculated from their relationships with  $w_m$ .

Equation (20) can also be transformed into the following Volterra integral equation:

$$w_m(T) - w_m(T - \eta)H(T - \eta) + \int_0^T G_m(\tau)w_m(T - 2\eta - \tau)H(T - 2\eta - \tau)d\tau = w_m^0(T - \eta)H(T - \eta). \quad (25)$$

On close examination, however, the solution of Eq. (25) requires repeating exactly the same convolutions as in Eq. (24). Therefore it is quite unnecessary to use the numerical techniques of integral equations, even if only one single integral equation is involved, for obtaining the inverse Laplace transform of Eq. (20). On the contrary, the use of two simultaneous Volterra integral equations for obtaining the inverse Laplace transforms of  $\bar{w}_m$  and  $\bar{w}_m^e$ , as in Ref. 6 would have unduly increased the numerical difficulty and the computation effort.

### Results and Discussions


Numerical results are obtained for a case in which both shells are made of steel and both fluids are water. The material properties and dimensions used are such that

$$\begin{aligned} c_r = \rho_r = 1, \quad C^2 = C_e^2 = 17.79133 \\ M^e = 32.09875, \quad M = 6.41975 \\ h^e/a^e = 1/250, \quad h/a = 1/50 \\ \zeta = 0.8, \quad \eta = 0.2. \end{aligned} \quad (26)$$


The incidence is a step wave with  $f(s) = 1/s$ . The transient responses of the shells are calculated for four transit times of the incident wavefront, i.e.,  $T = 0$  to 8. For this duration of time, it requires twenty successive convolution integrations in Eq. (24). The parabolic rule is used for the numerical integration in conjunction with Trauboth's fast convolution scheme. Since numerical roundoff and truncation errors accumulate at each successive integration, the integration intervals  $\Delta\tau$  are kept sufficiently small to minimize these errors. The roots of  $A_m(s)$ , and from these,  $w_m^0(T)$ , and  $G_m(T)$ , are evaluated with high accuracy before starting the convolution integration. It is also found that, similar to the case of a single shell [7], eight terms ( $m = 0 - 7$ ) in the series solution are sufficient for the representations of the shell deflections, velocities, and stresses. The integration intervals used are:  $\Delta\tau = 0.01$  for  $w_0$  and  $w_1$ ,  $\Delta\tau = 0.0025$  for  $w_2$ , and  $\Delta\tau = 0.00125$  for the higher terms. If the solution must be carried out for shorter time duration, say up to  $T = 4$ , a much coarser  $\Delta\tau$  will suffice.

In all subsequent figures, the present solutions are plotted in dotted lines and compared to the solutions obtained by the method of Ref. 7 and plotted in solid lines for the case in which the outer shell is absent.

Figure 2 shows the results for the  $w_0$ 's and  $\dot{w}_1$ 's. Similar to the single-shell case, the present results approach their respective asymptotic values after about two transit times, and

Fig. 2 - Time histories of  $w_0$  and  $w_1$ 

the numerical asymptotes of  $w_0$  and  $w_1$  agree with formulae (17) and (19) respectively within three digits. These results are indicative of the accuracy of the present solution computation scheme. For this particular example, the outer shell is rather thin and its effect on the response of the inner shell is observable in the  $w_0$ ,  $w_1$ , and  $w_2$  terms. Its effect on the higher modes of the inner shell, i.e.,  $w_m$  with  $m > 2$ , is quite undiscernible. Sample results for higher  $w_m$ 's are exhibited in Fig. 3. It can be seen there that the present results for  $w_3$  and  $w_4$  coalesce with the respective results for the outer shell absent case. The same can be said about  $w_5$ ,  $w_6$ , and  $w_7$ , and they are not replotted here.

Fig. 3 - Time histories of  $w_2$ ,  $w_3$ , and  $w_4$

Eight terms of  $w_m$ 's are summed in Eq. (7) for representing the total response of the inner shell. Figure 4 plots the time history of the relative radial deflection between the two apexes of the inner shell. The presence of the outer shell causes a downward shift of the curve representing the present result. This is due to the diminished  $w_0$  term.

Figure 5 shows the time histories of the radial velocities of the same two apexes. The presence of the outer shell causes a slight reduction of the velocities with little alteration of their profiles. Again this can also be expected by observing the slight reduction of the  $\dot{w}_1$  term.




Fig. 4 - Relative radial deflection between the apexes of the inner shell




Fig. 5 - Time histories of radial velocities at the apexes of the inner shell

It is reiterated here that the series solution in the form of Eq. (7) is not effective for obtaining the early time acceleration and pressure if the incident wave has a steep front. The customary remedy is to apply a modified Watson's transform to Eqs. (11) through (14) and calculate the asymptotic results for large  $s$  [1]. This is, however, much more complex than the single shell case studied in Ref. 12, and is outside the scope of the present report.

The polar membrane stress (force per unit length)  $N_\theta$  and the hoop membrane stress  $N_\phi$  are calculated by

$$N_\theta = \frac{aC^2}{(1 + \nu)M\zeta} \left[ (1 + \nu)w + \frac{\partial u}{\partial \theta} + \nu u \cot \theta \right] \quad (27)$$

and

$$N_\phi = \frac{aC^2}{(1 + \nu)M\zeta} \left[ (1 + \nu)w + \nu \frac{\partial u}{\partial \theta} + u \cot \theta \right]. \quad (28)$$



The time histories of the polar and/or hoop stress at  $\theta = \pi/2$  and at  $\theta = \pi$  are plotted in Fig. 6 and 7 respectively. The shielding of the outer shell reduces the stresses slightly. The mean stresses also decrease in proportion to  $w_0$ . The maximum stresses occur after the incident wave has completely engulfed the outer shell and are about twice the corresponding static values.




Fig. 6 - Time histories of the polar stress at  $\theta = \pi/2$




Fig. 7 - Time histories of the polar and hoop stresses at  $\theta = \pi$

All results of the present example infer that a thin outer shell tends to be transparent to the incident pulse.

In conclusion, it can be said that an accurate solution method for this problem has been developed and can be used for parametric studies. Moreover, a set of numerical results has been meticulously obtained to serve as a data base for the development of numerical methods for the analysis of the transient response of double hulls of practical configurations. One immediate scheme is to apply the currently available Doubly Asymptotic Technique [2] in the outer fluid field and the fluid finite element technique for the entrained fluid.

#### ACKNOWLEDGMENT

The work was supported by Defense Nuclear Agency under DNA subtask 99QAXSF502, Work Unit 10, Title Double Hull Response Evaluation, with SEA 03511 of Naval Sea Systems Command as the secondary manager. The author also wishes to express his appreciation to



Mr. G.J. O'Hara, Lt. R. Wade, Dr. M. Baron, and Dr. T.L. Geers for their helpful discussions. Specifically, it was Dr. Geers who brought the author's attention to and made available the English translation of Ref. 6.

## REFERENCES

1. Überall, H., and Huang, H., "Acoustical Response of Submerged Elastic Structures Obtained Through Integral Transform," *Physical Acoustics*, Mason, W.P., and Thurston, R.N., eds., Academic Press, New York, 1977, pp. 217-275.
2. Geers, T.L. "Transient Responses Analysis of Submerged Structures," *Finite Element Analysis of Nonlinear Structure Behavior*, Belytachko, T., Osias, J.R. and Marcel. P.V., eds., ASME, AMD-14, 15-84 (1975).
3. Geers, T.L., "Excitation of an Elastic Cylindrical Shell by A Transient Acoustic Wave," *J. Appl. Mech.*, **35**, 459-469 (1969).
4. Huang, H., "A Qualitative Appraisal of the Double Asymptotic Approximation for Transient Analysis of Submerged Structures Excited by Weak Shock Waves," NRL Memorandum Report 3135, September 1975.
5. Huang, H., Everstine, G.C. and Wang, Y.F., "Retarded Potential Techniques for the Analysis of Submerged Structures Impinged by Weak Shock Waves," *Computational Methods for Fluid-Structure Interaction Problems*, Belytschko, T. and Geers, T.L., eds., ASME, AMD-26, 83-93 (1977).
6. Babaev, A.E. and Kubenko, V.D., "Interaction of a Nonstationary Pressure Wave with a System of two Spherical Shells," *Mekh. Tverdogo Tela*, **2**, 135-141 (1977).
7. Huang, H., "Transient Interaction of Plane Acoustic Waves with a Spherical Elastic Shell," *J. Acoust. Soc. Amer.*, **45**, 661-670 (1969).
8. Abramowitz, M. and Stegun, I.A., eds., *Handbook of Mathematical Functions*, AMS 55, National Bureau of Standard, Washington, D.C., 1964, Chap. 10, pp. 437-445.
9. Van der Pol, B. and Bremmer, H., *Operational Calculus*, Cambridge University Press, New York, 1964, Chap. VII, pp. 122.
10. Ibid, Chap. IX, p. 189.
11. Trauboth, H.H., "Recursive Formulae for the Evaluation of the Convolution Integral," *J. Assoc. Comput. Mach.* **16**, (No. 1), 63-72 (1969).
12. Tang, S. and Yen, D., "Interaction of a Plane Acoustic Wave with an Elastic Spherical Shell," *J. Acoust. Soc. Amer.* **47**, 1325-1333 (1970).

**DISTRIBUTION LIST**

**ASSISTANT TO THE SECRETARY DEFENSE  
ATOMIC ENERGY  
WASHINGTON , DC 20301  
ATTN DONALD COTTER**

**DIRECTOR  
DEFENSE ADVANCED RSCH PROJ AGENCY  
ARCHITECT BUILDING  
1400 WILSON BLVD  
ARLINGTON VA. 22209  
ATTN A TACHMINDJI  
ATTN STO KENT KRESA  
ATTN TECHNICAL LIBRARY  
ATTN R CHAPMAN**

**DIRECTOR  
DEFENSE INTELLIGENCE AGENCY  
WASHINGTON D.C. 20301  
ATTN DI-7D E. OFARRELL  
ATTN DI-7E  
ATTN DT-1C J. VERONA  
ATTN DT-2 WPNS + SYS DIV  
ATTN TECHNICAL LIBRARY**

**DIRECTOR  
DEFENSE NUCLEAR AGENCY  
WASHINGTON D.C. 20305  
ATTN STTL TECH LIBRARY  
ATTN STST ARCHIVES  
ATTN DDST  
ATTN SPSS**

**( 2 COPIES )**

**( 2 COPIES )**

**CHAIRMAN  
DEPT OF DEFENSE EXPLO SAFETY BOARD  
RM-GB270, FORRESTAL BUILDING  
WASHINGTON D.C. 20301  
ATTN DD/S+SS**

DIR OF DEFENSE RSCH + ENGINEERING  
WASHINGTON D.C. 20301

ATTN AD/SW  
ATTN DD/TWP  
ATTN DD/S+SS  
ATTN AD/NP

COMMANDER  
FIELD COMMAND  
DEFENSE NUCLEAR AGENCY  
KIRTLAND AFB, NM 87115  
ATTN FCTA  
ATTN FCTA-D

INTERSERVICE NUCLEAR WEAPONS SCHOOL  
KIRTLAND AFB, NM 87115  
ATTN TECH LIB

DIRECTOR  
JOINT STRAT TGT PLANNING STAFF JCS  
OFFUTT AFB  
OMAHA, NB 68113  
ATTN STINFO LIBRARY

WEAPONS SYSTEMS EVALUATION GROUP  
400 ARMY NAVY DRIVE  
ARLINGTON VA 22202  
ATTN DOC CON

CHEIF OF RES, DEV + ACQUISITION  
DEPARTMENT OF THE ARMY  
WASHINGTON D.C. 20310  
ATTN TECHNICAL LIBRARY  
ATTN DAMA-CSM-N LTC E. DEBOESER JR



COMMANDER  
HARRY DIAMOND LABORATORIES  
WASHINGTON D.C. 20438  
ATTN AMXDO-NP  
ATTN AMXDO-TI    TECH LIB

DIRECTOR  
U S ARMY BALLISTIC RESEARCH LABS  
ABERDEEN PROVING GROUND, MD 21005  
ATTN TECH LIB-E. BAICY

COMMANDER  
U S ARMY COMM COMMAND  
FORT HUACHUCA, AZ 85613  
ATTN TECHNICAL LIBRARY

COMMANDER  
U S ARMY MAT + MECHANICS RSCH CTR  
WATERTOWN, MA 02172  
ATTN R SHEA

COMMANDER  
U S ARMY NUCLEAR AGENCY  
FORT BLISS, TX 79916  
ATTN TECH LIB

COMMANDER  
U S ARMY WEAPONS COMMAND  
ROCK ISLAND ARSENAL  
ROCK ISLAND, IL 61201  
ATTN TECHNICAL LIBRARY

CHIEF OF NAVAL MATERIAL  
NAVY DEPT  
WASHINGTON D.C. 20360  
ATTN MAT 0323



CHIEF OF NAVAL OPERATIONS  
NAVY DEPARTMENT  
WASHINGTON D.C. 20350  
ATTN OP 03FG  
ATTN OP 985F

CHIEF OF NAVAL RESEARCH  
DEPARTMENT OF THE NAVY  
ARLINGTON VA 22217  
ATTN N PERRONE CD 474  
ATTN TECHNICAL LIBRARY

OFFICER IN CHARGE  
CIVIL ENGR LAB  
NAVAL CONST. BATTALION CTR  
PORT HUENEME CA 93041  
ATTN R ODELLO  
ATTN TECH LIB

COMMANDER  
NAVAL ELECTRONIC SYSTEMS COMMAND  
NAVAL ELECTRONIC SYSTEMS COMMAND HQS  
WASHINGTON D.C. 20360  
ATTN PME 117-21A

COMMANDER  
NAVAL FACILITIES ENGINEERING COMMAND  
HEADQUARTERS  
WASHINGTON D.C. 20390  
ATTN TECHNICAL LIBRARY

SUPERINTENDENT  
NAVAL POSTGRADUATE SCHOOL  
MONTEREY CA 93940  
ATTN CODE 2124 TECH RPTS LIBRARIAN

DIRECTOR  
NAVAL RESEARCH LABRATORY  
WASHINGTON D.C. 20375  
ATTN CODE 2027 TECHNICAL LIB  
ATTN CODE 840 J GREGORY  
ATTN CODE 8440 F ROSENTHAL  
ATTN CODE 8403A G OHARA  
ATTN CODE 8442 H HUANG

(3 COPIES)

(4 COPIES)

COMMANDER  
NAVAL SEA SYSTEMS COMMAND  
NAVY DEPARTMENT  
WASHINGTON D.C. 20362  
ATTN ORD - 91313 LIB  
ATTN CODE 03511 C POHLER

COMMANDER  
NAVAL SHIP RSCH AND DEVELOPMENT CTR  
UNDERWATER EXPLOSIONS RSCH DIVISION  
PORTSMOUTH V.A. 23709  
ATTN E PALMER  
ATTN TECHNICAL LIBRARY

COMMANDER  
NAVAL SHIP ENGINEERING CENTER  
CENTER BUILDING  
HYATTSVILLE MD 20782  
ATTN NSEC 6120D  
ATTN NSEC 6110.01  
ATTN NSEC 6105G  
ATTN NSEC 6105  
ATTN 6105C1  
ATTN TECHNICAL LIBRARY

COMMANDER  
NAVAL SHIP RESEARCH AND DEV CENTER  
BETHESDA MD 20034

ATTN CODE 17 WW MURRAY  
ATTN CODE 142-3 LIBRARY  
ATTN CODE 174  
ATTN CODE 11  
ATTN CODE 2740 Y WANG  
ATTN CODE 1962  
ATTN CODE 1903  
ATTN CODE 1731C  
ATTN CODE 1171  
ATTN CODE 19

COMMANDER  
NAVAL SURFACE WEAPONS CENTER  
WHITE OAK  
SILVER SPRING MD 20910

ATTN CODE 241 J PETES  
ATTN CODE 1224 NAVY NUC PRGMS OFF  
ATTN CODE 730 TECH LIB  
ATTN CODE 240 H SNAY  
ATTN CODE 243 G YOUNG

COMMANDER  
NAVAL SURFACE WEAPONS CENTER  
DAHLGREN LABORATORY  
DAHLGREN VA 22448  
ATTN TECHNICAL LIBRARY

COMMANDER  
NAVAL UNDERSEA CENTER  
SAN DIEGO, CA 92152  
ATTN TECHNICAL LIBRARY

COMMANDER  
NAVAL WEAPONS CENTER  
CHINA LAKE CA 93555  
ATTN CODE 533 TECH LIB



COMMANDING OFFICER  
NAVAL WEAPONS EVALUATION FACILITY  
KIRTLAND AIR FORCE BASE  
ALBUQUERQUE NM 87117  
ATTN TECHNICAL LIBRARY

DIRECTOR  
STRATEGIC SYSTEMS PROJECTS OFFICE  
NAVY DEPARTMENT  
WASHINGTON D.C. 20376  
ATTN NSP-272  
ATTN NSP-43 TECH LIB

AF CAMBRIDGE RSCH LABS, AFSC  
L.G. HANSCOM FIELD  
BEDFORD MA 01730  
ATTN SUOL AFCRL RSCH LIB

HEADQUARTERS  
AIR FORCE SYSTEMS COMMAND  
ANDREWS AFB  
WASHINGTON D.C. 20331  
ATTN TECHNICAL LIBRARY

COMMANDER  
ARMAMENT DEVELOPMENT & TEST CENTER  
ELGIN AFB FL 32542  
ATTN TECHNICAL LIBRARY

LOS ALAMOS SCIENTIFIC LAB  
P O BOX 1663  
LOS ALAMOS NM 87544  
ATTN DOC CONTROL FOR REPORTS LIB

SANDIA LABS  
LIVERMORE LAB  
P O BOX 969  
LIVERMORE CA 94550  
ATTN DOC CON FOR TECH LIB

SANDIA LABORATORIES  
P.O. BOX 5800  
ALBUQUERQUE NM 87115  
ATTN DOC CON FOR 3141 SANDIA RPT COLL

U S ENERGY RSCH & DEV ADMIN  
DIVISION OF HEADQUARTERS SERVICES  
LIBRARY BRANCH G-043  
WASHINGTON D C 20545  
ATTN DOC CONTROL FOR CLASS TECH LIB

UNIV OF CALIFORNIA  
LAWRENCE LIVERMORE LAB  
P.O. BOX 808  
LIVERMORE CA 94550  
ATTN TECHNICAL LIBRARY

AGBABIAN ASSOCIATES  
250 NORTH NASH STREET  
EL SUGONDO CA 90245  
ATTN M AGBABIAN

BATTELLE MEMORIAL INSTITUTE  
505 KING AVENUE  
COLUMBUS OH 43201  
ATTN TECHNICAL LIBRARY

BELL TELEPHONE LABORATORIES INC.  
MOUNTAIN AVE  
MURRAY HILL NJ 07974  
ATTN TECH RPT CTR

BOEING COMPANY  
P.O. BOX 3707  
SEATTLE WA 98124  
ATTN AEROSPACE LAB

CAMBRIDGE ACOUSTICAL ASSOC  
1033 MASSACHUSETTS AVE  
CAMBRIDGE MA 02138  
ATTN M JUNGER

CIVIL / NUCLEAR SYSTEMS CORP  
1200 UNIVERSITY N. F.  
ALBUQUERQUE NM 87102  
ATTN T DUFFY

ELECTRIC BOAT DIV  
GENERAL DYNAMICS CORP.  
GROTON CN 06340  
ATTN L. CHEN

GENERAL ELECTRIC CO.  
TEMPO-CENTER FOR ADVANCED STUDIES  
816 STATE STREET (P.O. DRAWER 00)  
SANTA BARBARA CA 93102  
ATTN DASAC

ITT RESEARCH INST  
10 WEST 35TH ST  
CHICAGO IL 60616  
ATTN TECHNICAL LIBRARY



INST FOR DEFENSE ANALYSIS  
400 ARMY NAVY DRIVE  
ARLINGTON VA 22202  
ATTN IDA LIBRARIAN R SMITH

J.L. MERRITT  
CONSULTING + SPECIAL ENGR SVS INC  
P.O. BOX 1206  
REDLANDS CA 92373  
ATTN TECHNICAL LIBRARY

KAMAN AVIDYNE  
DIV OF KAMAN SCEINCES CORP  
83 SECOND AVE  
NW INDUSTRIAL PARK  
BURLINGTON MA 01803  
ATTN E CRISCIONE  
ATTN TECHNICAL LIBRARY  
ATTN G ZARTARIAN

KAMAN SCIENCES CORP.  
P.O. BOX 7463  
COLORADO SPRINGS CO 80933  
ATTN TECHNICAL LIBRARY

LOCKHEED MISSILES AND SPACE CO.  
3251 HANOVER ST  
PALO ALTO CA 94304  
ATTN TECH INFO CTR D/COLL  
ATTN T GEERS D/52-33 BLDG 205

NATHAN M. NEWMARK  
CONSULTING ENGINEERING SERVICES  
1114 CIVIL ENGINEERING BLDG  
URBANA IL 61801  
ATTN N NEWMARK

POLYTECHNIC INST OF NEW YORK  
DEPT OF APPLIED MECH  
333 JAY STREET  
BROOKLYN NY 11201  
ATTN J KLOSNER

R+D ASSOCIATES  
P.O. BOX 3580  
SANTA MONICA CA 90403  
ATTN TECHNICAL LIBRARY

STANFORD RESEARCH INST  
333 RAVENSWOOD AVE  
MENLO PARK CA 94025  
ATTN SRT LIB ROOM G021  
ATTN B GASTEN  
ATTN G ABRAHAMSON

TETRA TECH INC.  
630 N ROSEMEAD BLVD  
PASEDNA CA 91107  
ATTN LI-SAN HWANG  
ATTN TECH LIB

THE BDM CORP  
1920 ALINE AVE  
VIENNA VA 22180  
ATTN TECH LIB

UNIV OF MARYLAND  
DEPT OF CIVIL ENGR  
COLLEGE PARK MD 20742  
ATTN B BERGER

URS RESEARCH CO.  
155 BOVET RD.  
SAN MATEO CA 94402  
ATTN TECH LIB

WEIDLINGER ASSOCIATES  
3000 SAND HILL ROAD  
BUILDING 4 SUITE 245  
MENLO PARK CA 94025  
ATTN J. ISENBERG

WEIDLINGER ASSOCIATES  
110 EAST 59TH STREET  
NEW YORK, NY 10022  
ATTN DR. M. BARON

ASSISTANT CHIEF FOR TECHNOLOGY  
OFFICE OF NAVAL RESEARCH  
ARLINGTON, VIRGINIA 22217  
ATTN CODE 200

TELEDYNE BROWN ENG.  
MAIL STOP 44  
300 SPARKMAN DRIVE  
RESEARCH PARK  
HUNTSVILLE, ALABAMA 35807  
ATTN DR. MANU PATEL

DIRECTOR  
U S ARMY WATERWAYS EXPERIMENT STN  
P.O. BOX 631  
VICKSBURG MS 39180  
ATTN J STRANGE  
ATTN W FLATHAU  
ATTN TECH LIB (UNCL ONLY)



AF INSTITUTE OF TECHNOLOGY, AU  
WRIGHT PATTERSON AFB, OH 45433  
ATTN LIB AFIT BLDG 640 AREA B (UNCL ONLY)

NKF ENGINEERING ASSOCIATES, INC.  
8720 GEORGIA AVENUE  
SILVER SPRING, MD. 20910  
ATTN DR. R.O. BELSHEIM

M&T CO.  
2130 ARCH STREET  
PHILADELPHIA, PA. 19103  
ATTN B. McNAIGHT

CATHOLIC UNIV. OF AMERICA  
DEPT OF PHYSICS AND INSTITUTE  
OF OCEAN SCIENCE AND ENGINEERING  
WASHINGTON, D.C.  
ATTN H. ÜBERALL

CORNELL UNIV.  
COLLEGE OF ENGINEERING  
DEPT OF THEORETICAL AND  
APPLIED MECHANICS  
ITHACA, N.Y. 14850  
ATTN Y.H. PAO

SANDIA LABORATORIES  
SANDIA CORPORATION  
DIVISION 9324  
ALBUQUERQUE, NEW MEXICO 87115  
ATTN M.J. FORRESTAL